

Performance of multivoltine mulberry silkworm (*Bombyx Mori* Linn.) topically treated with the phytojuvenoid hormone

* Roli Srivastava, VB Upadhyay

Silkworm laboratory, Department of Zoology, D.D.U. Gorakhpur University, Gorakhpur, Uttar Pradesh, India

Abstract

The phytojuvenoid activates the physiological and developmental processes in the living beings; therefore, the topical application of phytojuvenoid on *Bombyx mori* larvae has been studied to observe any possibility for the heavy production of silk. The experiments were conducted with the phytojuvenoid concentration of 10, 20, 30 and 40 % and number of treatment viz; single, double, triple. The total RNA content in the haemolymph of *Bombyx mori* larvae at the initial and final stage of spinning was significantly ($P < 0.01$) influenced due to variation in the phytojuvenoid concentration but the number of larval treatment did not significantly influence the RNA content. Maximum level of RNA content was recorded in 30% phytojuvenoid concentration – triple treated larvae. The phytojuvenoid influences the level of protein in the larvae and caused some beneficial effect on the life pattern of silkworm and the productivity of cocoon.

Keywords: phytojuvenoid, RNA content, haemolymph, stage of larvae, *bombyx mori*

1. Introduction

Sericulture, or silk farming, is the rearing of silkworms for the production of silk. Although there are several commercial species of silkworms, *Bombyx mori* is the most widely used and intensively studied silkworm. Nistari is a resistant variety of multivoltine mulberry silkworm (*Bombyx mori*) which contributes up to a great extent in the commercial production of cocoon. In order to increase, the production of silk, efforts have been made to study effect of ecological factor^[1], photoperiod^[2] etc on the performance of silkworm. The Magnetization of eggs influences silk producing potential^[3] and incubation period of eggs^[4] and larval performance^[5]. The phytoecdysteroid has been noticed to influence the development, growth, silk producing and reproductive potential of *B. mori*^[6, 7, 8, 9, 10]. The juvenile hormone analogue also has been noticed to influence the reproductive potential and biochemical constituents of *Bombyx mori*^[11, 12, 13]. The synchronized maturation of larvae and simultaneous spinning of cocoon is very important in the sericulture industry. However, the response to such treatment varies depending on the dosage of compounds showing duration and number of applications^[14]. The more food ingested during this period gets converted and it turn contributes to silk protein. Delay in moulting is probably due to the inhibitory action of JH on ecdysone synthesis in *B. mori*^[15]. JH is claimed to inhibit protein synthesis in early treated larvae with later on region protein synthesis resulting in bigger silk gland and the result is improvement of cocoon shell weight^[16]. In the present study *Pinus longifolia* was taken for experiment due to its good availability and containing juvenile compound. This study has academic as well as economic importance in the field of sericulture.

2. Material and Method

The seed cocoons (pupa enclosed in silken case) of multivoltine mulberry silkworm (*Bombyx mori* nistari) were obtained from the silkworm grainage, Directorate of sericulture, Behraich Uttar

Pradesh and were maintained in the plywood trays (23 x 20 x 5cm) under the ideal rearing conditions^[16] in the silkworm laboratory, Department of Zoology D.D.U. Gorakhpur University, Gorakhpur. The temperature and relative humidity were maintained at $26 \pm 1^\circ\text{C}$ and $80 \pm 5\%$ RH respectively till the emergence of moths from the seed cocoons. The newly emerged moths were quickly picked up and kept sex-wise in separate trays to avoid copulation. The male moths were smaller in size but more active than the female moths which were comparatively larger and less active. The whole grainage operation was performed as per description given by Krishnaswamy *et al.* (1973)^[17]. Moths have a tendency to pair immediately after emergence, therefore, Sufficient pairs, each containing one male and one female from newly emerged moths were allowed to mate at $26 \pm 1^\circ\text{C}$ and $80 \pm 5\%$ RH in 12 hour / day dim light condition. After four hours of mating, the paired moths were detached manually. The female moths were allowed for egg laying. After 24 hours of egg laying, the female moths were individually examined for their disease freeness and after formaline treatment the dried eggs were transferred to the incubator for hatching. After hatching, the silkworm larvae were reared on the fresh and clean leaves of *Morus alba* given as food in the rearing trays.

These larvae were taken for the purpose of experiments. After completion of fifth instar, the ripe worms ceased feeding and ready for spinning. Small mountages were provided to the ripe worms and thus, sufficient number of cocoons was obtained from the silkworm larvae reared in our laboratory.

2.1 Design of Experiment

For extraction of phytojuvenoid the needle of *Pinus* were collected, washed thoroughly with distilled water and dried in incubator at 37°C . The dried materials were powdered separately with the help of mechanical device. Further, 50 gm powder was subjected to extraction separately through soxlet apparatus with 250 ml distilled water for 40 hours. After 40 hours of extraction

a little amount of concentrated solution of plant extract was obtained. The concentrated solution was dried and 6.45 gm material was obtained in powdered form. The dried powder thus obtained, was dissolved in distilled water as 5 gm in 25 ml water and used this solution for further experiment, as 100% concentration of phytojuvenoid. For further experiment the suitable narrow ranges of *Pinus* phytojuvenoid concentrations viz. 10, 20, 30 and 40% were taken. Thus, four phytojuvenoid concentrations were applied topically by spraying as 1 ml on to 100 larvae separately. Three sets of experiments were designed viz., single, double and triple treatment of larvae.

2.2 Single treatment of larvae

Single treatment of larvae was performed at the initial stage of fifth instar larvae just after fourth moulting. One hundred larvae of fifth instar at the initial stage were taken out from the BOD incubator and treated with one ml of 10% concentrated solution of *Pinus* needle extract by sprayer.

2.3 Double treatment of larvae

Double treatment of larvae was started from the initial stage of fourth instar larvae. In the first treatment, one hundred larvae of fourth instar were treated by 1 ml of 10% concentrated solution of *Pinus* needle extract by spraying. The treated larvae were then transferred in BOD incubator for rearing and development. Further, similar second treatment for the same larvae was given at the initial stage of fifth instar larvae. Thus, in double treatment, fourth and fifth instar larvae were treated.

2.4 Triple treatment larvae

For triple treatment, the third instar larvae in the initial stage were separated from BOD incubator. In the first treatment one hundred, third instar larvae, were treated by 1 ml of 10% concentrated solution of *Pinus* needle extract by sprayer and kept in BOD for rearing. The second treatment of same larvae was done just after third moulting i. e. at the initial stage of fourth instar larvae and transferred in BOD incubator for rearing. Third treatment was given at the initial stage of fifth instar i.e. just after fourth moulting of the same treated larvae as earlier. Thus, in the triple treatment third, fourth and fifth instar larvae were treated.

Similar experiments were performed by 20, 30 and 40% concentrations of phytojuvenoid obtained from *Pinus* needle extract. A control set was always maintained with each set of experiment.

To observe the effect of phytojuvenoid at various stages of *Bombyx mori* larvae on total RNA contents in the haemolymph of larvae at the initial and final stage of spinning, following method was adopted. The estimation of RNA was performed according to Schneider (1957) [18] by using the orcinol reagents. For measuring the RNA content took 0.62 mg (0.5 ml) haemolymph from the Vth instar larval stage and pupae. The tissues, thus obtained were homogenized separately in 5% TCA and centrifuged at 5000 rpm for 20 minutes. Took one ml of supernatant and added 2ml of distilled water and 3.0 ml of orcinol reagent (1 gm orcinol, 100ml concentrated HCl and 0.5 gm Ferric Chloride). The reaction mixture was kept in boiling water-bath for 20 minutes. The greenish blue-colour developed was measured at 660nm. Standard curves were drawn using different concentrations of yeast RNA as standard. RNA has been expressed as µg/mg tissue. Six replicates of each experiment were made.

Similarly, other series of experiments were performed with silkworm larvae, treated with 20, 30 and 40% phytojuvenoid concentrations separately with a set of control. All the data obtained were analyzed statistically by two-way ANOVA and Post- hoc test.

3. Results

3.1 Total RNA content in the haemolymph of larvae at the initial stage of spinning

The data given in table-1a clearly indicates that the phytojuvenoid concentration and number of larval treatment influenced the total RNA content in the haemolymph of the larvae at the initial stage of spinning. With the increasing number of larval treatment with 10, 20 and 30% phytojuvenoid concentration, the total RNA content in the haemolymph of the larvae at the initial stage of spinning increased gradually and reached to the maximum level of 2.52 ± 0.05 µg/mg in case of triple treated larvae with 30% phytojuvenoid concentration. In case of larval treatment with 40% phytojuvenoid concentration, the total RNA content in the haemolymph of larvae at the initial stage of spinning increased in single treated larvae but further increase in the number of larval treatment caused decline in the total RNA content in the haemolymph of the larvae at the initial stage of spinning which reached to the minimum level of 2.10 ± 0.03 µg/mg in triple treated larvae. The trend of increase in the total RNA content in the haemolymph of the larvae at the initial stage of spinning was almost same in 10, 20 and 30% phytojuvenoid concentration in relation to the number of larval treatment.

Two-way ANOVA indicates that variation in the phytojuvenoid concentration significantly ($P_1 < 0.01$) influenced the total RNA content but number of larval treatment did not cause significant influence on the total RNA content in the haemolymph of larvae at the initial stage of spinning. The Post-hoc test (table-1b) shows significant group difference in the total RNA content in between all the group combinations except in between control and 40%, 10 and 20% and 20 and 30% in single and double treatment of larvae. In triple treated larvae significant group difference in the total RNA content was recorded all the group combinations except in between 10 and 20% and 20 and 40% phytojuvenoid concentration.

3.2 Total RNA content in the haemolymph of larvae at the final stage of spinning

It is clear from the data given in the table-2a that the phytojuvenoid concentration and number of larval treatment influenced the total RNA content in the haemolymph of larvae at the final stage of spinning. With the increasing number of larval treatment with 10, 20 and 30% phytojuvenoid concentration, the total RNA content in the haemolymph of larvae at the final stage of spinning increased gradually and reached to the maximum level of 1.85 ± 0.07 µg/mg in case of triple treated larvae with 30% phytojuvenoid concentration. In case of the larval treatment with 40% phytojuvenoid concentration, the total RNA content in the haemolymph of larvae at the final stage of spinning increased in single treated larvae but further increase in the number of larval treatment caused decline in the total RNA content in the haemolymph of larvae at the final stage of spinning which reached to the minimum level of 1.56 ± 0.05 µg/mg in triple treated larvae. The trend of increase in the total RNA content in the haemolymph of larvae at the final stage of spinning was almost same in 10, 20

and 30% phytojuvenoid concentration in relation to the number of larval treatment.

Two-way ANOVA indicates that variation in the phytojuvenoid concentration significantly ($P_1 < 0.01$) influenced the total RNA content while number of larval treatment did not cause significant influence on the total RNA content in the haemolymph of the larvae at the final stage of spinning. The Post-hoc test (table-2b) shows significant group difference in the total RNA content in the haemolymph of the larvae at the

final stage of spinning in between control and 10%, control and 20%, 10 and 30%, 20 and 40% and 30 and 40% in single treated larvae. In the double treated larvae significant group difference in the total RNA content was noticed in between all the group combinations except in between 10 and 20%, 10 and 30% and 30 and 40% phytojuvenoid concentration. In triple treated larvae significant group difference in the total RNA content was noticed in all the group combinations except in between 10 and 20% and 20 and 30% phytojuvenoid concentration.

Table 1a: Effect of phytojuvenoid treatment on the total RNA content ($\mu\text{g}/\text{mg}$) in the haemolymph of *Bombyx mori* larvae at the initial stage of spinning

Stage of treatment (Larval instar)	Phytojuvenoid concentration (%)					F1-ratio n1=4
	Control X1	10 X2	20 X3	30 X4	40 X5	
Single	2.20	2.33	2.38	2.42	2.26	17.76*
(V)	± 0.04	± 0.03	± 0.02	± 0.05	± 0.07	
	2.20	2.37	2.42	2.47	2.18	
Double						
(IV-V)	± 0.04	± 0.05	± 0.06	± 0.03	± 0.04	
Triple	2.20	2.41	2.46	2.52	2.10	
(III-V)	± 0.04	± 0.08	± 0.07	± 0.05	± 0.03	

F2-ratio = 0.1724** n2=2 *P1<0.01 ** Non significant

Each value represents mean \pm S.E. of six replicates
X1, X2, X3, X4 and X5 are the mean values of the total RNA content ($\mu\text{g}/\text{mg}$) in the haemolymph in control, 10, 20, 30 and 40 % phytojuvenoid concentration respectively.

Table 1b: Post - hoc test showing effect of phytojuvenoid treatment on the total RNA content ($\mu\text{g}/\text{mg}$) in the haemolymph of *Bombyx mori* larvae at the initial stage of spinning.

Mean difference stage of treatment in between			
groups	Single	Double	Triple
X1~X2	*0.13	*0.17	*0.21
X1~X3	*0.18	*0.22	*0.26
X1~X4	*0.22	*0.27	*0.32
X1~X5	0.06	0.02	*0.10
X2~X3	0.05	0.05	0.05
X2~X4	*0.09	*0.10	*0.11
X2~X5	*0.07	*0.19	*0.31
X3~X4	0.04	0.05	0.06
X3~X5	*0.12	*0.24	*0.36
X4~X5	*0.16	*0.29	*0.42

$$\text{Honesty Significant difference (HSD)} = q \sqrt{\frac{\text{MS within}}{n}}$$

$$= 6.10 \sqrt{\frac{0.003}{6}}$$

$$= 0.07$$

MS=Mean square value of ANOVA table

q = studentized range static

n = No. of replicates

* = shows significant group difference

X1, X2, X3, X4 and X5 are the mean values of the total RNA content ($\mu\text{g}/\text{mg}$) in the haemolymph of *Bombyx mori* in control, 10, 20, 30 and 40 per cent phytojuvenoid concentration respectively.

Table 2a: Effect of phytojuvenoid treatment on the total RNA content ($\mu\text{g}/\text{mg}$) in the haemolymph of *Bombyx mori* larvae at the final stage of spinning.

Stage of treatment (Larval instar)	Phytojuvenoid concentration (%)					F1-ratio n1=4
	Control X1	10 X2	20 X3	30 X4	40 X5	
Single	1.62	1.67	1.72	1.77	1.66	14.61*
(V)	± 0.04	± 0.03	± 0.08	± 0.03	± 0.05	
Double	1.62	1.71	1.76	1.81	1.60	
(IV-V)	± 0.04	± 0.04	± 0.06	± 0.08	± 0.07	
Triple	1.62	1.75	1.80	1.85	1.56	
(III-V)	± 0.04	± 0.06	± 0.09	± 0.07	± 0.05	

F2-ratio = 0.6218** n2=2 *P1<0.01 ** Non significant

Each value represents mean \pm S.E. of six replicates
X1, X2, X3, X4 and X5 are the mean values of the total RNA content ($\mu\text{g}/\text{mg}$) in the haemolymph in control, 10, 20, 30 and 40 % phytojuvenoid concentration respectively.

Table 2b: Post - hoc test showing effect of phytojuvenoid treatment on the RNA content in the haemolymph of *Bombyx mori* larvae at final stage of spinning.

Mean difference stage of treatment in between			
groups	single	double	triple
X1~X3	*0.10	*0.14	*0.18
X1~X4	*0.15	*0.19	*0.23
X1~X5	0.04	0.02	*0.06
X2~X3	0.05	0.05	0.05
X2~X4	*0.10	*0.10	*0.10
X2~X5	0.01	*0.10	*0.19
X3~X4	0.05	0.05	0.05
X3~X5	*0.06	*0.16	*0.26
X4~X5	*0.11	*0.21	*0.29

$$\begin{aligned} \text{Honesty Significant difference (HSD)} &= q\sqrt{\frac{\text{MS within}}{n}} \\ &= 6.10\sqrt{\frac{0.02}{6}} \\ &= 0.06 \end{aligned}$$

MS = Mean square value of ANOVA Table

q = studentized range static

n = No. of replicates

* = shows significant group difference X_1, X_2, X_3, X_4 and X_5 are mean values of RNA content in the haemolymph of *Bombyx mori* larvae in control, 10, 20, 30 and 40 per cent phytojuvenoid concentration respectively.

4. Discussion

The level of RNA content in the haemolymph of *Bombyx mori* at the initial and final stage of spinning influenced due to variation in the concentration of phytojuvenoid and number of larval treatment of larvae. The RNA content was increased with the from 10 to 30% phytojuvenoid concentration while in 40% phytojuvenoid concentration it declined with the same number of treatment of larvae. The unique increase in RNA: DNA ratio in the haemolymph on the first day of spinning is correlated with the start of the degradation of nucleic acid during spinning period [19], whereas, the alteration in RNA/DNA ratio is indicative of a compensatory augmentation of net protein synthesis during the cold acclimation [20]. The nucleic acid content (DNA and RNA) was increased with the increasing magnetic field exposure of silkworm larvae till 3500 gauss while further increase in the strength of magnetic field caused notable decline in the nucleic acid content [21]. Comparative biometric studies have attempted to identify which of the silk gland parameters is the target; four parameters show high correlation with silk productivity of different strain of *B.mori*. They are number of silk gland cells, silk gland weight, the DNA and the RNA content [22]. Methoprene and fenoxycarb enhanced the fibroin and sericine which activate the silk gland to synthesize more DNA and RNA [23]. It is well known that the resistance to biotic and abiotic constraints is governed by polygene with complex inheritance patterns and with lot of environmental influences [24]. Thus, it is concluded that the larvae treated with 30% phytojuvenoid concentration at triple treatment is most suitable for the improvement of different economic characters and good status of different biochemical parameters in *Bombyx mori*. Phytojuvenoid enhanced the larval duration and the larvae consume more mulberry leaves, leading to an increase in the fibroin and sricine present in the food causing stimulatory effects on the silk gland. The higher concentration of the phytojuvenoid seems to cause stress response causing decline in the nucleic acids content.

Reference

1. Upadhyay VB, Gaur KP, Gupta SK. Effect of ecological factors on the silk producing potential of the mulberry silkworm (*Bombyx mori* Linn.). Malays. Appl. Biol. 2004; 33(1):13-18.
2. Mishra AB, Upadhyay VB. Nutritional efficiency of bivoltine *Bombyx mori* Linn. Larvae at different photoperiod. Proc. 80 session Indian Science Congress Assoc. Goa., 1993; 54-55.
3. Tripathi SK. Protien level changes under magnetic exposure of larvae in *Bombyx mori*: A multivoltine mulberry silkworm. Academic j. of Entomology. 2012; 5(2):73-80.
4. Upadhyay VB, Prasad S. Biotechnological importance of cocoon magnetization with particular reference to the reproductive potential of multivoltine mulberry silkworm (*Bombyx mori* Linn). Sericologia, 2010; 50(4):461-472.
5. Prasad S, Upadhyay VB. Biotechnological importance of cocoon magnetization with particular reference to the larval performance of multivoltine mulberry silkworm (*Bombyx mori* Linn.). Middle-East J. Sci. Res. 2011; 10(5):565-572.
6. Pandey P, Upadhyay VB. Phytoecdysteroid treatment on the larval performance of multivoltine mulberry Silkworm *Bombyx mori* Linn. Malays. Appl. Biol. 2013; 42(1):51-60.
7. Pandey P, Upadhyay VB. Influence of phytoecdysteroid on silk producing potential of Multivoltine mulberry silkworm (*Bombyx mori* Linn.). J. Appl. Biosci., 2012; 38(2):187-191.
8. Pandey P, Upadhyay VB. Phytoecdysteroid treatment on the larval performance of multivoltine mulberry Silkworm *Bombyx mori* Linn. Malays. Appl. Biol., 2013; 42(1):51-60.
9. Srivastava K, Upadhyay VB. Influence of Phytoecdysteroid on Weight of Ovary and Weight of Multivoltine Mulberry Silkworm (*Bombyx mori* Linn.). European Journal of Applied Sciences. 2012; 4(3):117-122.
10. Srivastava K, Upadhyay VB. Effect of Phytoecdysteroid on Length of Silk Filament and Non-Breakable Filament Length of Multivoltine Mulberry Silkworm *Bombyx mori* Linn. Academic Journal of Entomology, 2012b; 5(3):174-181.
11. Srivastava R, Upadhyay VB. Effect of Bioactive Phytojuvenoid on the Reproductive Potential of Multivoltine Mulberry Silkworm *Bombyx mori* Linn. (Lepidoptera: Bombycidae). Academic Journal of Entomology 2013; 6(2):85-92.
12. Srivastava R, VB Upadhyay. Application of phytojuvenoid enhances the protein level in the multivoltine mulberry silkworm (*Bombyx mori* Linn.) International Journal of Fauna and Biological Studies; 2014; 1(3):21-26.
13. Srivastava R, Upadhyay VB. Biochemical constituents of multivoltine mulberry silkworm (*Bombyx mori* Linn.) Influenced by Phytojuvenoid compound. International Journal of Fauna and Biological Studies; 2016; 3(1):01-05.
14. Chowdhary SK, Raju PS, Ogra RK. Effects of JH analogues on silkworm, *Bombyxmori* L. growth and development of silk gland. Sericologia, 1990; 30(2):155-165.
15. Trivedy K, Nair KS, Assan NM. Datta RK. A Juvenile hormone mimic modulated enhancement of silk productivity in silkworm, *Bombyx mori* Indian J. Serc., 1997; 360:35-38.
16. Garel JP. The physiology and biology of spinning in *Bombyx mori* V Endocrinological aspects of silk production. Experientia, 1983; 39(5):461-466.
17. Krishnaswamy S, Narasimhanna MM, Suryanaryan SK, Raja S, Kumar. Sericulture Manual 2. Silkworm Rearing. F.A.O. Agric. Serves Bull. Rome. 1973; 15(2):1-131.
18. Schneider WC. Determination of nucleic acids in the tissue of pentose analysis. In "Methods in Engymology" (eds., S.P ccolmick and N.D.Kaplan) Academic Press, 1957; 14:680-681.
19. Okabe K, Koyanagi R, Kogo K, RNA in the degenerating silk gland of *Bombyx mori* J. Insect Physiol. 1975; 21(6):1305-1310.
20. Hazel JR, Prosser CL. Biochemistry of different tissue in

- Philosamia ricini*. Adv. Insect Physiol. London, 1974; 1: 111-174.
21. Chaugale AK, Pawar BK, NK More. Influence of the magnetic field on the nucleic acids in developing silk gland of *Bombyx mori*. Indian J. Seric, 1995; 34(2):161-164.
 22. Fayard JM. Production de soie et caractéristiques des glands sericigènes de 13 races de ver à soie (*Bombyx mori*). Ann. Genet. Sel. Anim., 1978; 10:259-270.
 23. Mamatha DM, Cohly HPP, Raju AHH, Rajeswara M. Rao. Studies on the quantitative and qualitative characters of cocoons and silk from methoprene and fenoxycarb treated *Bombyx mori* (L.) larvae. African Journal of Biotechnology, 2006; 5(15):1422-1426.
 24. Promboon A, Shimada T, Fujiwara H, Kobayashi M. Linkage map of random amplified polymorphic DNAs (RAPDs) in the silkworm, *Bombyx mori*. Gen. Res. 1995; 66:1-7.