



## Induced breeding of *Labeo mrigala* through single dose of ovaprim-C at carp seed farm, Alsu, district Mandi, Himachal Pradesh

Rakesh Kumar

Department of Biotechnology and Life Sciences, Shoolini University, Himachal Pradesh, India

### Abstract

A study was conducted to observe the effect of intramuscular injection of ovaprim-C on the number of eggs/kg, fertilization rate and hatching percentage at Carp Seed Farm, Alsu, District Mandi (Himachal Pradesh), during July to August 2022, on *Labeo mrigala* (Mori). Studied fish specimens were spawned successfully following a single dose of injection of ovaprim-C with 0.4 ml kg<sup>-1</sup> for female and 0.05 ml kg<sup>-1</sup> for male brooders. Ova and milt were stripped simultaneously and mixture was stirred for 20 to 35 s during which fertilization occurred. Hatching occurred within 25 to 40 h after fertilization. The experiment was conducted in circular Chinese hatchery with 4 m diameter. Overall fertilization and hatchling% age was 65.00% and 75.125% respectively. Wet body weight was observed to have a positive influence on absolute ( $r=0.983$ ) and relative fecundity ( $r = 0.910$ ) in log-log scale.

**Keywords:** *Labeo mrigala*, ovaprim-c, male brooders, hormone injection

### Introduction

The experiments were conducted on males and females of a major Indian carp Mrigal, *Labeo mrigala*, by injecting ovaprim during induced breeding. Induced breeding is a method in which exogenous hormones are injected into the body of mature parent fish for induction of breeding (Heggberget, 1996) [6]. Marte *et al.* (1987) [15] reported that a drug known as ovaprim has been commonly used as a spawning hormone in fish breeding. Induced breeding through hypophysation has been the common practice since the development of the technology in 1957, several synthetic commercial formulations of purified salmon gonadotropin and dopamine antagonists such as ovaprim, ovatide and wova-FH have also been successfully used in recent years (Jena *et al.*, 1998) [9]. Ovaprim is a commercial product used as a spawning aid in fishes and contains a salmon gonadotropin-releasing hormone analog and a dopamine antagonist. Ovaprim has been shown to be effective and safe in numerous ornamental finfish species in the refereed scientific literature. Many years of research investigating hormonal triggers for ovulation and spermiation went into the development of Ovaprim (Peter *et al.*, 1986) [28]. Human chorionic gonadotropin (HCG) has been used to induce final maturation of oocytes and also as a tool for utilization in commercial aquaculture (Kelly and Kohler, 1994; Mylonas *et al.*, 2009) [11, 17]. The use of different forms of gonadotropin releasing hormone agonist (GnRH<sub>a</sub>), which stimulate secretion of endogenous gonadotropin (GTH) (Zohar, 1989; Zohar and Mylonas, 2001) [32, 33], ovaprim and ovatide are a kind of analogue of salmon gonadotropin releasing hormone (sGnRH<sub>a</sub>) with a dopamine blocker (Syndel International Inc., 2003) [31]. The use of sGnRH<sub>a</sub> resulted in successful stimulation of ovulation in some cyprinids (Drori *et al.*, 1994; Hill *et al.*, 2005) [5, 7] and catfishes (Sahoo *et al.*, 2007) [30]. The objective of induced ovulation is to produce, on demand, a large supply of high quality eggs. Egg quality is assessed by characteristics such as egg fertility and hatching (Bromage

and Cumaranatunga, 1988) [2]. The aims and objectives of this study were to assess the fecundity, that is, number of egg/kg of wet body weight, by artificial spawning in *L. mrigala*.

### Materials and methods

The experiments were conducted at Carp Seed Farm, Alsu, District Mandi (Himachal Pradesh), during July to August 2022, on *Labeo mrigala* (Mori). 15 specimens of *L. mrigala* were used for investigation, they weighed about 1.0 to 2.0 kg. These artificial breeding tests were conducted in circular chinese spawning hatchery with 4 m in a diameter with optimum water quality variables. Ovaprim-C was injected in a single dose. These experiments were done to produce fish fry for sale for farmers and achieve targets by the following procedures.

### Selection and handling of bloodstocks

Good quality and well-matured male and female brood fish were collected because according to Muir and Robert (1985) [16], healthy parental fish are prerequisite for successful artificial propagation. The female fish with a bulging abdomen and swollen soft, pink red genital papilla were selected. The males that released milt when subjected to gentle pressure on the abdomen were selected. The sex ratios of one female to two males were used, spawning run so as to get best results.

### Hormone injection

The brood fish were weighed and dosage of Ovaprim-C solution was calculated according to Nandeesh *et al.* (1991) [23] by the following formula:

Quantity to be injected (ml) = weight of brood fish (kg) x dosage of ovaprim-C (ml)

The males were injected ovaprim-C with about 0.05 ml/kg, while the females were injected ovaprim-C at a rate of 0.4 ml/kg. It was ensured that all the equipments were cleaned and needle of the hypodermic syringe was cleaned by cotton swab, soaked in alcohol before injection. Required amount of Ovaprim-C was withdrawn from the bottle by keeping the

needle upward and syringe was squeezed gently to expel any trapped air. During injection, a brood fish was placed in a cloth bag, lying laterally in water, the upper half of the fish was held above the water surface. At the inner side of the basal part of the pectoral fin, where it was scale less, the syringe needle was inserted gently towards the head at an angle of 45° to the body's longitudinal axis to a depth of about 1.5 cm and fluid was injected slowly. Then, both males and females were released in the fresh aerated water of circular tank.

### Estrus

After injection, the spawning behavior was observed in the brood fish as the males started chasing the females after 8:30 h. This phenomenon is called estrus in which the female shows restlessness and its abdomen and tail becomes extremely constricted. The activity lasted for about 30 to 60 min after which fish were netted out for stripping.

### Stripping

For spawning, the belly of the female was pressed. The eggs were released from the genital pore and were collected in the dry plastic bowl. The same process was repeated with the males and milt was put into the bowl having eggs.

### Fertilization

Following the semi-dry fertilization method, milt was mixed with the eggs using a bird feather for two minutes. The

significance of this method was that, the travel distance for the sperm to the micropyle of the egg was short. Then eggs were washed with water for 10 min, they absorbed water and attained the size of 1 to 1.5 mm in diameter.

### Assessment of fecundity

The total number of eggs spawned was counted by the volumetric method. In this method, approximately 1 g egg sample was weighed out three times. The total number of eggs spawned was calculated by multiplying the average number of eggs from three one gram samples with the total weight of eggs sampled. So, the total number of eggs/kg was counted by the following formula:

**Table 1:** Effect of Ovaprim-C on spawning of *L. mrigala*.

Parameter	Ovaprim treatment
No. of females treated.	15
Total weight of females	35.00 kg
Total no. of eggs	4000000
Total no. of fertilized eggs	2560000
Total no. of hatchling	2000000
Overall fertilization percentage	64%
Overall hatching percentage	78.125%
Average no. of eggs/kg	60000
Average no. of fertilized eggs/kg	50000
Average no. of hatching/kg	39992

Number of eggs/kg = Total number of eggs/total weight of fish

**Table 2:** Spawning response of female *L. mrigala*

Month	Temperature (°C)	Number of female	Total weight of females (kg)	Dose of Ovaprim (ml kg-1)	Total Number of egg	Total Fertilization rate	Total number of hatching
June 2022	26	5	12	0.4	0.70	0.60	0.320
July 2022	26.4	5	11	0.4	0.80	0.65	0.297
August 2022	27	5	13	0.4	0.75	0.68	0.323

### Assessment of fertilization

After 3 to 4 h of fertilization, the fertilized and unfertilized eggs were distinguished. In the fertilized eggs, the division of cells was so regular that the size and shape of blastomeres was equal. So, the percentage of fertilized eggs was calculated according to Muir and Robert (1985) [16]; the percentage of eggs beyond morula stage expressed as percentage of the total number of eggs was the fertilization rate, using the following formula:

Percentage of fertilization = (Number of fertilized egg/total number of eggs) x 100

### Incubation and hatching

As the breeding tests were conducted in the circular spawning tanks with 2 m diameter, hatching occurred after 18 to 24 h at water temperature (20 to 24.5°C). The newly hatched larvae were kept in the circular spawning tanks with bolting cloth for three days until yolk was absorbed. Then the percentage of hatchling was calculated by the following formula:

Percentage of hatchlings = (Total number of hatchlings/total number of fertilized eggs) x 100

### Results

Mean values and ranges of absolute fecundity, fertilization and hatchlings of Rohu (*L. rohita*) following the intramuscular injection of ovaprim-C are given in Table 1. Fish given single injection of Ovaprim-C was successfully

induced to spawn. 15 females were injected with ovaprim-C. Spawning response of female *L. mrigala* is given in Table 2. Ovulation of fish in this treatment was 100%. Overall fertilization and hatchling% age was 64.00 and 78.125%, respectively (Table 1). Regression analyses of absolute and relative fecundity with body weight were found highly correlated ( $P < 0.001$ ).

### Discussion

In the present study, a single intramuscular injection of synthetic hormone, ovaprim-C resulted in successful spawning of *Labeo rohita*. The results of the current work in the hormonal stimulation are similar to the effectiveness and usefulness by using Ovaprim-C (Jamroz *et al.*, 2008) [8]. Although, according to Basaran and Sabsun (2008) [1], speed and gentleness during fish capture and handling are of utmost importance. Different analogues of LH-RH, without pituitary gland, which of results in failure of spawning indicates that dopamine blocks the action of LH-RH on the secretion gonadotropin reported by Naeem and Salam (2005) [18]. However, the use of dopamine antagonists like pimozide or doperidon, potentiate the action of LH-RH, resulting in successful spawning (Chang and Peter, 1983) [3]. Major breakthrough in the history of aquaculture happened when extensive research on Chinese carp (Peter *et al.*, 1988) [29] and a new Linpe method was used in which LH-RH analogue is combined with a dopamine antagonist. Then, Canada introduced the ovaprim-C containing the analogue

of salmon gonadotropin releasing hormone (D-Arg6, Pro9, Net) and dopamine antagonist, and studies conducted in India (Nandeeshia *et al.*, 1990a) <sup>[26]</sup> and Pakistan (Khan *et al.*, 1992; Naeem and Salam, 2005; Naeem *et al.*, 2005a; b) <sup>[12, 18, 20]</sup> revealed the superiority of Ovaprim-C in induced spawning. Dose of Ovaprim-C used in *L. mrigala* in the present experiment is 0.4 ml/kg, while experiments conducted by Nandeeshia *et al.* (1990a) <sup>[26]</sup> and Khan *et al.* (1992) <sup>[12]</sup> reported the dose rate of 0.3 to 0.4, 0.4 ml/kg, respectively. So, the present study reveals that Ovaprim-C use is more economical in commercial carp seed production, as it saves a considerable amount of time and avoids the excessive handling of brood fish. The positive response of *Mori* to ovaprim indicated the higher potency of this drug in inducing the spawning. *L. mrigala* have been reported to spawn with 10 mg of Pimozide (Kaul and Rishi, 1986) <sup>[10]</sup> and the results obtained from Ovaprim is supported by the results of the trials in 1988 and 1989 (Nandeeshia *et al.*, 1990a; b) <sup>[26]</sup>.

## References

- Basaran F, Sabsun N. Survival rates of black sea turbot (*Psetta maximamaeotica*.1.1758) broodstock captured by gill nets from different depths and their adaptation culture conditions. *Aqua. Int.*,2008;12:321-331.
- Bromage NR, Cumararatunga PRT. Egg production in rainbow trout. In: Muir JF, Roberts RJ (eds) Recent Advances in Aquaculture, Groom Helm Ltd., London, 1988, 65-138.
- Chang JP, Peter RE. Effect of Pimozide and des-Gly (D-Ala6) lutinising hormone releasing hormone, Ethyl amide on serum gonadotropin concentration, germinal vesical migration and ovulation in female gold fish *Carassius auratus*. *Gen. Comp. Endocrinol.*,1983;52:30-37.
- Crim LW, Glebe BD. Advancement and synchrony of ovulation in Atlantic salmon with pelleted LHRH analog. *Aquaculture*,1984;43:47-56.
- Drori S, Ofir M, Sivan BL, Yaron Z. Spawning induction in common carp (*Cyprinus carpio*) using pituitary extract or GnRH superactive analogue combined with methoclopramide: analysis of hormone profile, progress of oocyte maturation and dependence on temperature. *Aquaculture*,1994;119:393-407.
- Heggberget TG. The role of aquaculture in world fisheries. Oxford & IBH Publishing co. Pvt. Ltd. New Delhi, 1996.
- Hill JE, Baldwin JD, Graves JS, Leonard R, Powell GFF, Wanton CA. Preliminary observations of topical gill application of reproductive hormones for induced spawning of a tropical ornamental fish. *North American J. Aquaculture*,2005;67:7-9.
- Jamroz M, Kucharezky D, Hakuc Blazowska A, Krejszef S, Kujawa R, Kupren K. Used in the controlled production of IDE, LEUCISCUS IDUS. *Arch. Polish Fish.*,2008;16:363-370.
- Jena JK, Aravindakshan PK, Chandra S, Muduli HK, Ayyappan S. Comparative evaluation of growth and survival of Indian major carps and exotic carps in raising fingerlings. *J. Aquacult. Trop.*,1998;13:143-150.
- Kaula M, Rishi KK. Induced spawning of indian major carp *Cirrhinus mrigala* with LH-RH analogues or pimozide. *Aquaculture*,1986;54:45-48.
- Kelly AM, Kohler CC. Human chorionic gonadotropin injected in fish degrades metabolically and by cooking. *J. World Aquacult. Soc.*,1994;25:55-57.
- Khan MN, Janjua MY, Naeem M. Breeding of carps with Ovaprim (LH-RH Analogue) at Fish Hatchery Islamabad, Proceeding. Pakistan Congress of Zool.,1992;12:545-552.
- Lin HR, Peter RE. Hormones and spawning in fish. *Asian Fisheries Sci.*,1996;9:21-23.
- Lam TJ. Environmental influences on gonadal activity in fish. In: Fish Physiology. Hoar WS, Randall DJ, Donaldson EM (eds). Academic Press, London, 1983, 65-116.
- Marte CL, Sherwood NM, Crim LW, Harvey B. Induced spawning of maturing milkfish (*Chanos chanos forsskal*) with gonadotropin-releasing hormone (GnRH) analogues administered in various ways. *Aquaculture*,1987;60:303-310.
- Muir FJ, Robert JR. Recent advances in Aquaculture, Groom Helm, London and Sydney, Westview press Boulder, Calarado, 1985, 2.
- Mylonas CC, Fostier A, Zanuy S. Broodstock management and hormonal manipulation of fish reproduction. *Aquaculture*,2009;197:99-136.
- Naeem M, Salam A. Induced spawning of bighead carp *Aristichthys nobilis* by using Ovaprim-Cat fish hatchery Islamabad, Pakistan. *Sindh Univ. Res. J. (Sci. Ser.)*,2005;37(1):9-16.
- Naeem M, Salam A, Jafar A. Induced spawning of major carp *Catla catla* by a single intermuscular injection of Ovaprim-C and fecundity at fish hatchery Islamabad, Pakistan. *J. Biol. Sci.*,2005b;5(6):776- 780.
- Naeem M, Salam A, Diba F, Saghir A. Fecundity and Induced Spawning of Silver carp, *Hypophthalmichthys molitrix* by using a Single Intramuscular Injection of ovaprim-C at Fish Hatchery Islamabad, Pakistan, Pakistan. *J. Bio. Sci.*,2005a;8(8):1126-1130.
- Naeem M, Salam A, Ali M, Mehreen M, Ali M, Khan MJ, *et al.* Breeding performance of sustainable fish *Ctenopharyngodon idella* through single intramuscular injection of Ovaprim-C at Bahawalpur, Pakistan. *Afr. J. Biotechnol.*,2011b;10(57):12315-12318.
- Naeem M, Salam A, Elahi N, Ali M, Ishtiaq A, Andleeb A. Effect of body weight on absolute and relative fecundity of *Hypophthalmichthys molitrix* with intramuscular injection of Ovaprim - C. *Int. J. Agric. Biol.*,2011a;13:141-144.
- Nandeeshia MC, DAS SK, Ramacharya Varghes TJ. Further observations on breeding of carps with Ovaprim. *Spl. Publ. 6, AFSIB Mangalore*, 1991, 41.
- Nandeeshia MC, Keshavanath P, Varghese TJ, Shetty HPC, Rao KG. Alternate inducing agents for carp breeding, Progress in research. In: Keshavanath P, Radhakrishnan KV (eds) *Carp seed production technology*, 1990b.
- Proceedings of the workshop on carp seed production technology. 2-4 September. Special Publication 2. Asian fisheries society, Indian branch, Mangalore, India, 1988, 12-16.
- Nandeeshia MC, Rao KG, Jayanna R, Parker NC, TJ Varghese TJ, Keshavanath P, *et al.* Induced spawning of Indian major carps through single application of Ovaprim. In: Hirano R, Hanyu M (eds). *The second*

- asian fisheries forum, Asian Fisheries Society, Manila, Philippines, 1990a, 581-585.
27. Park IS, Kim HB, Choi HJ, Lee YD, Kang HW. Artificial induction of spawning by human chorionic gonadotropin (HCG) or carp pituitary extract (CPE) in olive flounder, *Paralichthys olivaceus*. *J. Aquacult*,1994;7:89-96.
  28. Peter R, Sokolowska M, Nahorniak CS. Comparison of (DArg6, trp7 Leu8 Pro9, Net) - luteinizing hormone (LHRH-A), in combination with pimozoide, in stimulating gonadotropin release and ovulation in the goldfish, *Carassius auratus*. *Can. J. Zool*,1986;65:987-991.
  29. Peter RE, Lin HR, Van der Kraak. Induced ovulation and spawning of cultured freshwater fish in China: Advances in application of GnRH analogues and dopamine antagonists. *Aquaculture*,1988;74:1-10.
  30. Sahoo SK, Giri SS, Chandra S, Sahu AK. Effect of Ovaprim doses and latency period on induced spawning of *Clarias batrachus*: Observation on larval deformity. *Indian J. Experiment Biol*,2007;45:920- 922.
  31. Syndel International Inc. Using Ovaprim to induce spawning in cultured fish, 2003.
  32. Zohar Y. Fish reproduction, its physiology and artificial manipulation. In: Shilo MC, Sargi SH (eds), *Fish culture in warm Water System, Problems and Trends*. CRC Press, 1989, 65-119.
  33. Zohar Y, Mylonas CC. Endocrine manipulation of spawning induction in cultured fish from hormone to gene. *Aquaculture*,2001;197:99-139.