



Morphometric differentiation of *Apis Cerana Indica* Across Distinct Ecological Zones in Nagpur, Maharashtra

Bahurupi Puja¹, Shinkhede Milind²

¹ Department of Zoology, RTMNU, Nagpur, Maharashtra, India

² Department of Zoology, Dada Ramchand Bakhru Sindhu Mahavidyalaya, Nagpur, Maharashtra, India

Abstract

Present study investigates the influence of distinct ecological zones agricultural and urban on the morphometric characteristics of *Apis cerana indica* population and field biology. A total of 120 adult bees, including drones, nurse bees, and foragers, were sampled from both habitats. Ten morphometric traits were measured, encompassing body length, head length, thorax and abdomen dimensions, antennal length, wing lengths, and leg lengths. The data reveal that bees from agricultural areas exhibited slightly larger dimensions across all measured traits compared to their urban counterparts, indicating a possible correlation between habitat quality and bee development. Floral diversity and seasonal pollen analysis further confirmed differential foraging patterns, with agricultural bees visiting a broader spectrum of nectar and pollen sources. These findings underscore the role of environmental factors including floral availability, landscape composition, and climate in shaping the physical development of *Apis cerana indica*, and highlight the importance of habitat conservation in ensuring sustainable pollinator health.

Keywords: *Apis cerana indica*, morphometric differentiation, pollen variability, agriculture vs. urban zones, nagpur

Introduction

Apis cerana indica, often known as 'Satpuda bees' in Maharashtra, serves as an efficient pollinator of regional crops and native plants. Globally, bees contribute to the pollination of over 70% of agricultural crops, with approximately 15% of major cultivated crops relying on domesticated bees for successful pollination (Abrol, 2012) [2]. Due to its adaptability and ecological significance, extensive research has been conducted on this species to evaluate its behavior, ecological preferences, and potential for sustainable apiculture (Abrol, 2009) [1]. The Nagpur district in central India features a variety of agro-climatic zones, urban areas, and industrial regions, supporting both cultivated and wild plant species.

Previous morphometric studies of *Apis cerana* have covered diverse ecological zones across India, from the Himalayas to the southern tip, including the Andaman Islands. Significant contributions include research by Rahman and Singh (1948) [20], Kapil (1956) [14], Deodikar (1962) [9], and Narayanan *et al.* (1961) [17]. However, limited literature addresses seasonal or regional morphometric variation (Alpatov, 1927; Ruttner *et al.*, 1978) [21]. The present study investigates the effects of environmental and geographical differences on morphometric traits of *Apis cerana indica* in agricultural and urban zones of Nagpur.

Pollen analysis from bee colonies serves as an indicator of floral preferences during different seasons (Bilisik *et al.*, 2008) [7]. Urbanization, habitat fragmentation, and pesticide use are major threats to bee diversity (Spivak *et al.*, 2011; Goulson *et al.*, 2005; Thompson, 2003; Gels *et al.*, 2002; Higes *et al.*, 2008; Benaets *et al.*, 2017) [5, 11, 12, 13, 22, 23]. This, studies suggest that sustainable practices such as organic farming can positively influence bee foraging behavior and colony health even in urban environments (Czech *et al.*, 2000) [8]. This study aims to elucidate the extent to which regional differences in flora and climatic conditions affect the development and physical characteristics of *Apis cerana indica*.

Materials and methods

The present study was carried out to see how differences in pollen and environmental conditions affect the physical traits of *Apis cerana indica* (F.). Two very different sites in Nagpur district, Maharashtra, were chosen for the colonies: an agricultural area in Karanbhad, Parseoni (Latitude: 21.362226, Longitude: 79.106343) and an urban area in Shankar Nagar, Nagpur (Latitude: 21.132188, Longitude: 79.056513).

Pollen samples were collected directly from the hive entrance to identify the floral sources visited by the bees. These samples were preserved in 70% ethanol and analyzed microscopically for floral origin characterization.

A total of 120 adult bees were taken for study, with 10 individuals from each of the three castes drones, nurse bees, and foragers collected from colonies at both sites. The bees were first anesthetized with chloroform and then placed in warm water so their body parts could be fully extended. After this, they were preserved in 70% ethanol for later study. All dissections and measurements were done under controlled laboratory conditions. Ten body characters were measured: body length, head length, thorax length, abdomen length, forewing length, hindwing length, front leg length, middle leg length, and hind leg length. Body length was measured with a digital Vernier caliper, while the other parts were measured with a stereomicroscope, using the method of Mattu and Verma (1984) [15]. The data collected were calculated and presented as mean, standard deviation (SD), and range of variation (RV).

Results and discussion

Tables 1 and 2 present the morphometric data for *Apis cerana indica* collected from different geographical regions, specifically agricultural and urban zones. These areas are characterized by distinct climatic conditions, including variations in temperature, humidity, and the surrounding flora and fauna (Michener, 1974; Velthuis, 1992) [16, 24]. The

morphological traits of *Apis cerana* were observed to vary with geographical location (Bhatta & Kumar, 2021) [6], indicating the influence of environmental conditions on the species' development.

Bees from agricultural areas were generally larger than those from urban regions across all castes. Drone, nurse, and worker bees in the agricultural zones had slightly bigger heads, thoraxes, and abdomens, suggesting better overall growth. Their antennae were also longer, which may help them sense and locate flowers more efficiently. Wings, both forewings and hindwings, were consistently longer in agricultural bees, particularly in workers, supporting longer and more effective flights during foraging. Legs of bees from farmland were longer as well, likely allowing them to move more easily and collect nectar and pollen from a wider

variety of plants. Among all castes, worker bees showed the most noticeable differences, reflecting their demanding role in foraging and maintaining the colony. These patterns indicate that the environmental conditions in agricultural areas such as richer floral diversity, greater food availability, and more favorable microclimates may contribute to better morphometric development compared to urban habitats. Overall, these findings highlight how habitat quality can directly influence the physical traits and potential performance of *Apis cerana indica*.

All morphometric values observed in this study were cross-checked with previous studies (Ananda, 2000; Patil & Pastagia, 2014; Pal *et al.*, 2008; Deodikar, 1962; Kapil, 1956) [4, 9, 14, 18, 19] to ensure accuracy.

Table 1: Quantitative data on significant morphological characters of the three castes of the Indian honey bee, *Apis cerana indica* (F.), collected from an agricultural area

Body characters	Drone		Nurse Bee		Forager Bees	
	Mean±SD	RV	Mean±SD	RV	Mean±SD	RV
Body Length	11.99±0.35	1.10	11.55±0.55	1.50	13.14±0.18	0.60
Head Length	3.40±0.14	0.40	2.92±0.16	0.50	3.50±0.14	0.40
Thorax Length	4.14±0.15	0.40	3.72±0.24	0.80	4.11±0.16	0.50
Abdomen Length	5.50±0.12	0.40	4.18±0.12	0.40	6.20±0.12	0.40
Antenna Length	4.99±0.10	0.30	4.56±0.11	0.30	4.85±0.11	0.30
Fore wings Length	8.73±0.13	0.40	7.75±0.13	0.30	9.71±0.13	0.40
Hind wings Length	6.24±0.10	0.30	5.49±0.12	0.40	6.51±0.09	0.20
Front legs Length	7.44±0.14	0.50	6.20±0.11	0.30	7.53±0.14	0.50
Middle legs Length	8.06±0.13	0.40	7.17±0.09	0.30	8.07±0.13	0.40
Hind legs Length	8.60±0.18	0.60	8.47±0.13	0.40	9.11±0.16	0.50

Table 2: Quantitative data on significant morphological characters of the three castes of the Indian honey bee, *Apis cerana indica* (F.), collected from an urban area

Body characters	Drone		Nurse Bee		Forager Bees	
	Mean±SD	RV	Mean±SD	RV	Mean±SD	RV
Body Length	11.81±0.37	1.10	11.11±0.22	0.60	12.25±0.45	1.30
Head Length	3.30±0.12	0.40	2.78±0.12	0.30	3.11±0.16	0.50
Thorax Length	3.97±0.14	0.30	3.12±0.14	0.40	3.33±0.13	0.40
Abdomen Length	4.09±0.12	0.40	3.67±0.09	0.30	4.55±0.11	0.30
Antenna Length	4.26±0.14	0.50	3.37±0.09	0.30	3.60±0.12	0.40
Fore wings Length	8.41±0.16	0.50	7.66±0.23	0.80	8.98±0.18	0.50
Hind wings Length	5.51±0.17	0.60	4.47±0.13	0.40	6.33±0.13	0.40
Front legs Length	6.47±0.09	0.30	6.07±0.16	0.50	6.85±0.13	0.40
Middle legs Length	7.40±0.12	0.40	6.31±0.16	0.50	7.70±0.12	0.40
Hind legs Length	8.50±0.12	0.40	8.00±0.12	0.40	8.94±0.11	0.30

SD- Standard Deviation, RV- Range of Variation All the mean values are in mm

Agricultural Zone

In the agricultural area, forager bees exhibited a wide spectrum of floral visitation, suggesting a diverse and seasonally changing floral landscape. During the winter months (January–February), the dominant pollen sources were *Alternanthera sessilis*, *Brassica campestris*, *Lagascea mollis*, and *Azadirachta indica*. Notably, *Azadirachta indica* contributed significantly in February (48.2%), indicating early blooming and high attractiveness due to its abundant nectar and pollen.

The spring months (March–April) reflected a shift in floral preference toward *Citrus sinensis* and *Citrus limetta*, with these species collectively accounting for over 70% of the pollen frequency in April. Their dominance during this period may be attributed to mass flowering and their high

quality nectar, which are known to attract forager bees extensively.

In the post-monsoon period (October–December), *Lagascea mollis*, *Parthenium hysterophorus*, *Gossypium*, *Cajanus cajan*, and *Capsicum annum* emerged as prominent sources. The high frequency of *Gossypium* and *Cajanus cajan* (both ≥32% in November and December) suggests the importance of cultivated crops in sustaining bee populations during periods when wild flora are limited. Interestingly, *Parthenium hysterophorus*, typically regarded as an invasive weed, was regularly foraged in both October and November, indicating its ecological relevance as a fallback floral resource.

Urban Zone

In contrast, the urban site presented a relatively limited but consistent floral spectrum. In the early months (January–March), *Azadirachta indica* consistently dominated (ranging from 40.3% to 48.6%), likely due to its widespread

occurrence in urban green spaces and roadside plantations. The presence of *Alternanthera sessilis* and *Tridax procumbens* further complemented the pollen diversity in February and March.

During April, urban foraging patterns diversified significantly, with *Psidium guajava* (38.1%), *Delonix regia*, *Bombax ceiba*, *Poaceae*, and *Cassia fistula* being visited. These ornamental and native species are commonly planted along roads and in parks, providing a transitional source of nectar and pollen between major flowering seasons.

In the post-monsoon period, *Alternanthera sessilis* remained a dominant floral source in October and November, while *Ocimum sanctum* and *Parthenium hysterophorus* also made substantial contributions. In December, *Albizia lebeck* (45.9%) and *Poaceae* (39.5%) were the major floral resources, showing the significance of late-flowering tree species and grasses in sustaining urban bee foraging.

Conclusion

The present study demonstrates that the morphometric characteristics of *Apis cerana indica* are significantly influenced by environmental variables, particularly the floral diversity and habitat type. Populations from agricultural regions exhibited relatively larger body sizes

and more developed morphological traits across all castes indicative of favorable ecological conditions such as diverse floral availability, consistent nectar sources, and relatively stable microclimatic parameters. These results are consistent with previous findings (Free, 1970; Rodionov & Shabarshov, 1986) ^[10, 21], supporting the notion that *Apis cerana* displays notable phenotypic plasticity in response to both biotic (vegetation type, resource availability) and abiotic (temperature, habitat structure) factors.

Comparative analysis of floral foraging patterns further reinforces this observation. Forager bees in agricultural zones accessed a broader and more seasonally varied array of floral resources, including both cultivated and wild flora. Conversely, urban colonies were limited to a narrower but steady set of ornamental and tree species, reflecting the constrained ecological offerings of urban landscapes.

Collectively, these findings highlight the importance of habitat quality and landscape heterogeneity in shaping both the morphology and foraging ecology of *Apis cerana indica*. Conservation and management strategies aimed at sustaining pollinator populations should therefore prioritize habitat diversification and floral resource enrichment, particularly in rapidly urbanizing regions.

Table 3: Representation of floral resources visited by forager bees in the agricultural zone.

Month	Flora visited by bees	Frequency of pollen
January 2023	<i>Alternanthera sessilis</i>	28.6%
	<i>Brassica campestris</i>	19.4%
	<i>Lagascea mollis</i>	16.5%
	<i>Justicia procumbens</i>	15.8%
	<i>Ocimum basilicum</i>	5.9%
February 2023	<i>Sphaeranthus indicus</i>	13.8%
	<i>Alternanthera sessilis</i>	30.3%
	<i>Azadiracta indica</i>	48.2%
	<i>Brassica compestris</i>	14.2%
March 2023	<i>Ocimum basilicum</i>	7.3%
	<i>Alternanthera sessilis</i>	20.7%
	<i>Brassica sp.</i>	10.4%
	<i>Cipadessa baccifera</i>	5.9%
	<i>Citrus sinensis</i>	29.3%
	<i>Citrus limetta</i>	33.8%
	<i>Solanum lycopersicum</i>	1.9%
April 2023	<i>Solanum lycopersicum</i>	8.7%
	<i>Cipadessa baccifera</i>	12.8%
	<i>Citrus sinensis</i>	38.2%
	<i>Citrus limetta</i>	42.1%
October 2023	<i>Lagascea mollis</i>	41.2%
	<i>Parthenium hysterophorus</i>	30.5%
	<i>Sphaeranthus indicus</i>	12.9%
	<i>Alternanthera sessilis</i>	15.4%
November 2023	<i>Parthenium hysterophorus</i>	16.2%
	<i>Gossypium</i>	35.3%
	<i>Capsicum annum</i>	19.1%
	<i>Cajanus cajan</i>	29.4%
	<i>Lagascea mollis</i>	4.8%
December 2023	<i>Parthenium hysterophorus</i>	9.4%
	<i>Gossypium</i>	35.5%
	<i>Cajanus cajan</i>	32.4%
	<i>Justicia procumbens</i>	2.8%
	<i>Capsicum annum</i>	15.1%

Table 4: Representation of floral resources visited by forager bees in the urban zone.

Month	Flora visited by bees	Frequency of pollen
January 2023	<i>Azadiracta indica</i>	48.6%
	<i>Alternanthera sessilis</i>	39.2%
	<i>Lagascea mollis</i>	12.2%
February 2023	<i>Azadiracta indica</i>	40.3%
	<i>Brassica compestris</i>	26.3%
	<i>Tridax procumbens</i>	33.3%
March 2023	<i>Azadiracta indica</i>	41.5%
	<i>Psidium guajava</i>	37.6%
	<i>Prosopis julifera</i>	20.9%
April 2023	<i>Psidium guajava</i>	38.1%
	<i>Delonix regia</i>	20.5%
	<i>Bombax ceiba</i>	16.3%
	Poaceae	10.2%
	<i>Cassia fisfula</i>	14.9%
October 2023	<i>Alernanthera sessilis</i>	40.4%
	<i>Parthenium hysterophorus</i>	20.2%
	<i>Ocimum santum</i>	31.4%
November 2023	<i>Alernanthera sessilis</i>	38.9%
	<i>Parthenium hysterophorus</i>	30.5%
	<i>Ocimum santum</i>	30.6%
December 2023	<i>Albizia lebbeck</i>	45.9%
	Poaceae	39.5%
	<i>Chrysanthemum sp.</i>	14.6%

Reference

- Abrol DP. Distinguishing characters of the cavity nesting honeybees. In: Bees and beekeeping in India. Kalyani Publishers, Ludhiana, 2009, 80–81
- Abrol DP. Pollination biology, biodiversity conservation and agricultural production. Springer, Dordrecht, Heidelberg, London, New York, 2012.
- Alpatov WW. Biometrical studies on variation and races of the honeybee (*Apis mellifera* L.). Quarterly Review of Biology, 1929;4:1–58.
- Ananda MR. Studies on morphometric characters and behavioural traits of Indian honeybee, *Apis cerana indica* Fabricius from different locations of Karnataka. M.Sc. thesis (unpublished), University of Agricultural Sciences, Bangalore, 2000.
- Benaets K, Van Geystelen A, Cardoen D, De Smet L, de Graaf DC, Schoofs L, *et al.* Covert deformed wing virus infections have long-term deleterious effects on honeybee foraging and survival. Proceedings of the Royal Society B: Biological Sciences, 2017;284(1848):20162149.
- Bhatta VR, Kumar AN. Survey of bee friendly flowering plants and bee-plant interaction in an urban green space in Bengaluru, India. Journal of Environmental Biology, 2021;42(5):1330–1337.
- Bilisik A, Cakmak I, Bicakci A, Malyer H. Seasonal variation of collected pollen loads of honey bees (*Apis mellifera* L. anatoliaca). Grana, 2008;47(1):70–77.
- Czech B, Krausman PR, Devers PK. Economic associations among causes of species endangerment in the United States. BioScience, 2000;50(7):593–601.
- Deodikar GB. Research at apicultural laboratory. Indian Bee Journal, 1962;24:83–91.
- Free JB. The flower constancy of bumblebees. The Journal of Animal Ecology, 1970;39:395–402.
- Gels JA, Held DW, Potter DA. Hazards of insecticides to the bumble bees *Bombus impatiens* (Hymenoptera: Apidae) foraging on flowering white clover in turf. Journal of Economic Entomology, 2002;95(4):722–728.
- Goulson D, Hanley ME, Darvill B, Ellis JS, Knight ME. Causes of rarity in bumblebees. Biological Conservation, 2005;122(1):1–8.
- Higes M, Martín-Hernández R, Botías C, Bailón EG, González-Porto AV, Barrios L, *et al.* How natural infection by *Nosema ceranae* causes honeybee colony collapse. Environmental Microbiology, 2008;10(10):2659–2669.
- Kapil RP. Variations in the biometrical characters of the Indian honey bee, *Apis cerana* F. Indian Journal of Entomology, 1956;18:440–457.
- Mattu VK, Verma LR. Morphometric studies on the Indian honeybee, *Apis cerana indica* F. Effect of seasonal variations. Apidologie, 1984;15(1):63–74.
- Michener CD. The social behavior of the bees. Belknap Press of Harvard University, Cambridge (UK), 1974.
- Narayanan ES, Sharma PL, Phadke KG. Studies on biometry of the Indian bees: Tongue length and number of hooks on the hind wings of *Apis cerana* F. collected from Madras state. Indian Bee Journal, 1961;23:3–9.
- Pal S, Patnaik HP, Satapathy CR. Morphometric diversity in *Apis cerana indica* F. worker bees of Orissa. Journal of Plant Protection and Environment, 2008;5(2):42–51.
- Patil PN, Pastagia JJ. Morphometrics of workers of Indian honey bee, *Apis cerana* F. in Gujarat. An International e-Journal, 2014;3(2):148–153.
- Rahman KA, Singh S. Preliminary studies in the bionomics of *Apis cerana indica*. Indian Journal of Entomology, 1948;10(1):63–73.
- Rodionov VV, Shabarshov IA. The fascinating world of bees. Imported Publication, 1986.
- Ruttner F, Tassencourt L, Louveaux J. Biometrical-statistical analysis of the geographic variability of *Apis mellifera* L. I. Material and methods. Apidologie, 1978;9(4):363–381.
- Spivak M, Mader E, Vaughan M, Euliss Jr NH. The plight of the bees. Environmental Science and Technology, 2011;45:34–38.

24. Thompson HM. Behavioural effects of pesticides in bees—their potential for use in risk assessment. *Ecotoxicology*,2003;12:317–330.
25. Velthuis HHW. Pollen digestion and the evolution of sociality in bees. *Bee World*,1992;73(2):77–89.